

Trapped surfaces, horizons and exact solutions in higher dimensions

José M. M. Senovilla

Departamento de Física Teórica, Universidad del País Vasco, Apartado 644, 48080 Bilbao, Spain.

Abstract. A very simple criterion to ascertain if $(D - 2)$ -surfaces are *trapped* in arbitrary D -dimensional Lorentzian manifolds is given. The result is purely geometric, independent of the particular gravitational theory, of any field equations or of any other conditions. Many physical applications arise, a few shown here: a definition of general horizon, which reduces to the standard one in black holes/rings and other known cases; the classification of solutions with a $(D - 2)$ -dimensional abelian group of motions and the invariance of the trapping under simple dimensional reductions of the Kaluza-Klein/string/M-theory type. Finally, a stronger result involving *closed* trapped surfaces is presented. It provides in particular a simple sufficient condition for their absence.

PACS Numbers: 04.50.+h, 04.20.Cv, 04.20.Jb, 02.40.Ky

In 1965 Penrose [1] introduced in General Relativity (GR) the concept of *closed trapped surface*, which was crucial for the development of the singularity theorems and the study of gravitational collapse, black holes, cosmological expansion and several types of horizons, see e.g. [2, 3]. Trapped surfaces (closed or not) are 2-dimensional imbedded spatial surfaces such that any portion of them has, at least initially, a decreasing area along *any* future evolution direction. The term “closed” is used if the surfaces are compact without boundary [1, 2, 3].

This concept carries over to general Lorentzian manifolds $(\mathcal{V}, g)$ of any dimension D [4]. To fix ideas and notation, let S be a $(D - 2)$ -dimensional surface with intrinsic coordinates $\{\lambda^A\}$ ($A, B, \dots = 2, \dots, D - 1$) imbedded into the spacetime $\mathcal{V}$ by the parametric equations

$$x^\alpha = \Phi^\alpha(\lambda^A) \quad (\alpha, \beta \dots = 0, 1, \dots, D - 1). \quad (1)$$

S is alternatively locally defined by two independent relations $F_1(x^\mu) = 0$ and $F_2(x^\mu) = 0$. The tangent vectors $\{\vec{e}_A\}$ of S are

$$\vec{e}_A \equiv e_A^\mu \frac{\partial}{\partial x^\mu} \Big|_S \equiv \frac{\partial \Phi^\mu}{\partial \lambda^A} \frac{\partial}{\partial x^\mu} \Big|_S$$

so that the first fundamental form of S in $\mathcal{V}$ reads

$$\gamma_{AB} \equiv g_{\mu\nu} \Big|_S \frac{\partial \Phi^\mu}{\partial \lambda^A} \frac{\partial \Phi^\nu}{\partial \lambda^B} \quad (2)$$

which gives the scalar products of the $\{\vec{e}_A\}$ in $(\mathcal{V}, g)$. Assume that γ_{AB} is positive definite so that S is *spacelike*. Then, the two linearly independent normal one-forms $k_\mu^\pm$ to S can be chosen to be null and future directed everywhere on S , so they satisfy

$$k_\mu^\pm e_A^\mu = 0, \quad k_\mu^+ k^{+\mu} = 0, \quad k_\mu^- k^{-\mu} = 0, \quad k_\mu^+ k^{-\mu} = -1, \quad \text{on } S \quad (3)$$

where the last equality is a condition of normalization. Obviously, there still remains the freedom

$$k_\mu^+ \longrightarrow k'^+_\mu = \sigma^2 k_\mu^+, \quad k_\mu^- \longrightarrow k'^-_\mu = \sigma^{-2} k_\mu^- \quad (4)$$

where σ^2 is a positive function defined on S .

The two null (future) second fundamental forms of S are given by

$$K_{AB}^\pm \equiv -k_\mu^\pm e_A^\nu \nabla_\nu e_B^\mu \quad (5)$$

and their traces are

$$K^\pm \equiv \gamma^{AB} K_{AB}^\pm, \quad (6)$$

where γ^{AB} is the contravariant metric on S : $\gamma^{AC} \gamma_{CB} = \delta_B^A$. The scalar defining the trapping of S is then

$$\kappa \equiv 2 K^+ K^- = H_\mu H^\mu \quad (7)$$

where $\vec{H} \equiv -K^- \vec{k}^+ - K^+ \vec{k}^-$ is the mean curvature vector of S [4]. Clearly, κ and $\vec{H}$ are invariant under (4). S is said to be *trapped* (respectively *marginally trapped*, *absolutely non-trapped*) if κ is positive, (resp. zero, negative) everywhere on S . S is called *untrapped* otherwise. See [2] and section 4 in [3] for details and examples. Notice that S is trapped (resp. absolutely non-trapped) when $\vec{H}|_S$ is timelike (resp. spacelike). If $\vec{H}$ is null at a point, then at least one of the traces $K^\pm$ vanishes there, so that a necessary condition for S to be marginally trapped is that $\vec{H}|_S$ be null.

The meaning of the trapping is simple: the traces (6) are in fact the expansions of the two families of null geodesics emerging orthogonally from S , which are tangent to $\vec{k}^\pm$ at S [2, 3]. Thus, S is trapped if both null geodesics families are converging, or diverging, all over S .

In this letter, a very simple way to check the trapping of surfaces is found, and thereby a definition of horizon will also arise naturally. Without loss of generality, the family of $(D-2)$ -dimensional spacelike surfaces S_{X^a} can be described by $\{x^a = X^a\}$, with $a, b, \dots = 0, 1$, where X^a are arbitrary constants and $\{x^\alpha\}$ are local coordinates in $(\mathcal{V}, g)$. The line-element can be written as

$$ds^2 = g_{ab} dx^a dx^b + 2g_{aA} dx^a dx^A + g_{AB} dx^A dx^B \quad (8)$$

where $g_{\mu\nu}(x^\alpha)$ and $\det g_{AB} > 0$. There remains the freedom

$$x^a \longrightarrow x'^a = f^a(x^b), \quad x^A \longrightarrow x'^A = f^A(x^B, x^C) \quad (9)$$

keeping the form (8) and the chosen family of surfaces. Using coordinate conditions one can try to achieve $g_{aA} = 0$ or other similar simplifications, which are in fact useful in many applications, but I prefer to keep the full generality.

Let us calculate the scalar κ of (7) for the surfaces S_{X^a} . Clearly, the imbedding (1) for these surfaces is given locally by

$$x^a = X^a = \text{const.}, \quad x^A = \lambda^A,$$

and therefore the first fundamental form (2) for each S_{X^a} is $\gamma_{AB} = g_{AB}(X^a, \lambda^C)$. The null normals to each S_{X^a} satisfying (3) can be chosen as

$$\mathbf{k}^\pm = k_b^\pm dx^b|_{S_{X^a}}, \quad g^{ab}k_a^\pm k_b^\pm|_{S_{X^a}} = 0, \quad g^{ab}k_a^+ k_b^-|_{S_{X^a}} = -1 \quad (10)$$

(g^{ab} is not necessarily the inverse of g_{ab} !). Now the calculation of the $K_{AB}^\pm$ in (5) is straightforward:

$$K_{AB}^\pm = -k_c^\pm \Gamma_{AB}^c|_{S_{X^a}}$$

where $\Gamma_{\mu\nu}^\rho$ are the Christoffel symbols. From its definition $2\Gamma_{AB}^a = -g^{a\rho}(2\partial_{(A}g_{B)\rho} - \partial_\rho g_{AB})$ so that, by setting

$$G \equiv +\sqrt{\det g_{AB}} \equiv e^U, \quad \mathbf{g}_a \equiv g_{aA} dx^A \quad (11)$$

and using $k^{\pm B} = -\gamma^{BA} g_{aA} k^{\pm a}$, the two traces (6) can be obtained ($f_{,\mu} = \partial_\mu f$)

$$K_{X^a}^\pm = k^{\pm a} \left(\frac{G_{,a}}{G} - \frac{1}{G} (G \gamma^{AB} g_{aA})_{,B} \right) \Big|_{S_{X^a}}. \quad (12)$$

Hence, the mean curvature one-form reads

$$\boxed{H_\mu = \delta_\mu^a (U_{,a} - \text{div} \vec{g}_a)} \quad (13)$$

where div is the divergence operator on vectors at each S_{X^a} , so that the scalar (7) for each S_{X^a} is finally

$$\boxed{\kappa_{X^a} = -g^{bc} H_b H_c|_{S_{X^a}}} \quad (14)$$

(13-14) are the desired formulae, which are invariant under changes of type (4) and (9). Observe that one only needs to compute the norm of H_a as if it were a one-form in the 2-dimensional metric g^{ab} . The function $G = e^U$, which from (11) gives the canonical $(D-2)$ -volume element of the surfaces S_{X^a} (their area in $D = 4$), arises as a fundamental object. As (13) shows, $\mathbf{H}$ has a pure divergence term in general. However, as we are going to see presently, in many situations $\text{div} \vec{g}_a = 0$, in which case $H_a = U_{,a}$ and only the normal variation of volume is relevant. Let us stress that (13-14) are purely geometric, independent of any matter contents, of energy or causality conditions [2, 3], and of any field equations. They hold in general dimension D , including, in particular, the case of GR for $D = 4$.

In general, H_a will change its causal character at different regions. The hypersurface(s) of separation $\mathcal{H}$, defined locally by the vanishing of $g^{bc} H_b H_c$, is a fundamental place in $(\mathcal{V}, g)$ that I call the S_{X^a} -horizon. This contains (i) the regions with marginally trapped S_{X^a} , and (ii) the parts of each S_{X^a} where one of the traces vanishes. $\mathcal{H}$ coincides in many cases with the classical horizons, as shown in the examples that follow.

Many interesting applications can be derived from (13-14). Let us start with a simple illustrative example, the Kerr metric in Boyer-Lindquist coordinates $\{t, r, \theta, \phi\}$ (notation as in [2]). The case of physical interest arises for $\{x^a\} = \{t, r\}$, $\{x^A\} = \{\theta, \phi\}$. It is immediate to obtain $\mathbf{g}_r = 0$, $\mathbf{g}_t = 2amr\rho^{-2}\sin^2\theta d\phi$ and

$$e^{2U} = \sin^2\theta[(r^2 + a^2)\rho^2 + 2mra^2\sin^2\theta]$$

so $\text{div}\vec{g}_a = 0$, $H_a = U_{,a}$, and using (14) one easily derives (for $r > 0$) $\text{sign}\kappa_{t,r} = -\text{sign}\Delta$, with $\Delta = r^2 - 2mr + a^2$. This is the standard result, which identifies the classical event and Cauchy horizons at $\Delta = 0$ as well as the *closed* trapped surfaces at $\Delta < 0$.

Let us consider now the general spherically symmetric line-element in arbitrary D

$$ds^2 = g_{ab}(x^c)dx^a dx^b + R^2(x^c)d\Omega_{D-2}^2 \quad (15)$$

where $d\Omega_{D-2}^2$ is the round metric on the $(D-2)$ -sphere and $\det g_{ab} < 0$. Here $H_a = U_{,a} \propto R_{,a}/R$, and $\mathcal{H}$ is the classical apparent horizon [2, 3], which in particular becomes an event/Cauchy horizon in symmetric cases. The former case includes D -dimensional Robertson-Walker cosmologies, and the latter the Reissner-Nordström-Tangherlini black holes [5], among many others. A D -generalization of the standard “mass function” in spherical symmetry (see in GR, e.g., [6]) arises

$$2M(x^a) \equiv R^{D-3} (1 - g^{bc}R_{,b}R_{,c})$$

so that $2M(X^a) > R^{D-3}(X^a)$ for trapped $(D-2)$ -spheres S_{X^a} , which in this case are obviously closed. This agrees (up to a constant $(D-2)$ -volume factor) with [7].

For more up-to-date matters, let us apply the above results to the 5-dimensional *rotating black rings/holes* recently presented in [8] (containing a subset of the rotating black holes in [9]). Using the notation in [8] for the metric appearing in their formula (13), the physically relevant case arises for $\{x^a\} = \{\chi, y\}$, $\{x^A\} = \{v, x, \phi\}$. By simple inspection one reads off

$$\mathbf{g}_y = 0, \quad \mathbf{g}_\chi = \frac{\sqrt{v}(y - \xi_2)}{\sqrt{\xi_1}A} \frac{F(x)}{F(y)} dv, \quad e^{2U} = -\frac{F^3(y)}{A^4(x-y)^4}$$

and a very simple calculation gives $\text{div}\vec{g}_\chi = 0$ (ergo $H_a = U_{,a}$ once again) and, for the scalar (14)

$$\kappa_{\chi,y} = \frac{A^2(x-y)^2}{4F(x)F(y)} \left(\frac{3F'(y)}{F(y)} + \frac{4}{x-y} \right)^2 G(y).$$

Recalling that $F(y)F(x) < 0$ one gets $\text{sign}\kappa_{\chi,y} = -\text{sign}G(y)$ except at $\Sigma : y + 3x = 4\xi_1$, where $\kappa_{\chi,y} = 0$. This is due to the vanishing of $K_{\chi,y}^\pm$ at the *2-surfaces* $(\Sigma \cap S_{\chi,y}) \subset S_{\chi,y}$. One can check that Σ is located at $y > \xi_1$. Thus, there are *closed* trapped 3-surfaces $S_{\chi,y}$ for some $y \in (\xi_1, \xi_4)$ ($G(y)$ changes sign at ξ_4), while they are non-trapped for $y > \xi_4$. The horizon formed by closed marginally trapped surfaces is located at $y = \xi_4$, which is the event horizon described in [8]. In general, $\mathcal{H} = \Sigma \cup \{y = \xi_4\}$.

Another interesting application arises from the “generalized Weyl solutions” constructed recently in [10]. The main aim in [10] was to obtain the D -generalization

of the static and axisymmetric solutions of vacuum Einstein's equations. However, many other solutions not of Weyl-type were implicitly, maybe inadvertently, found. The general metric of [10] (for the “non-Weyl” case characterized by having real coordinates $\{Z, \bar{Z}\}$ in the notation of [10]) can be written in the form (8) by putting $g_{aA} = 0$, $g_{AB} = \text{diag}\{e^{2U_2}, \dots, e^{2U_{D-1}}\}$, and letting g_{ab} and $\{U_A\}$ to depend only on $\{x^a\}$. As proved with a particular coordinate choice in [10], the Ricci-flat condition for (8) implies then that $G = e^U$ satisfies (∇ is covariant derivative for g^{ab})

$$g^{ab}G_{;ab} = 0, \quad g^{ab}(G U_{A,a})_{;b} = 0 \quad \forall A. \quad (16)$$

These two expressions are conformally invariant with respect to g^{ab} . The first is simply the wave equation in the 2-metric g^{ab} for G , easily solvable in appropriate coordinates. The second relation (16) is identical with the equation in $D = 4$, so that in what follows one could always write down for each of the U_A the widely known solutions found in GR. Notice, though, that the proper choice of coordinates depends on the particular physical situation to be tackled. For instance, a simple possibility would be

$$g_{ab}dx^a dx^b = F^2(t, x)(-dt^2 + dx^2), \quad G = x. \quad (17)$$

However, from the previous analysis, this immediately implies that $\kappa_{t,x} < 0$, so that *all* the surfaces $t, x = \text{const.}$ are *absolutely non-trapped*.^Σ In other words, the choice (17) is adequate only for the regions of the spacetime with absolutely non-trapped $S_{t,x}$, and without $S_{t,x}$ -horizon $\mathcal{H}$. Analogous cases are given for instance by $G = \cosh t \sinh x$. These situations are appropriate to describe cylindrical or plane symmetric spacetimes, providing D -generalizations of this type of solutions in GR. This case is mentioned in [10]. There are, however, other physically inequivalent situations depending on the causal character of $U_{,a}$. These are essentially the following (keeping always the form of g_{ab} in (17) for simplicity):

1. $G = t$. Now all surfaces $S_{t,x}$ are trapped and again there is no $\mathcal{H}$. This case describes cosmological solutions, as for instance the D -dimensional Kasner metric [11], given by $U_A = p_A \log t$ with $\sum_A p_A = 1$. Analogous cases are given by $G = \sinh t \cosh x$.

2. $G = G(t - x)$ (or $G = G(t + x)$). In this case $\kappa_{t,x} = 0$ and all the surfaces $S_{t,x}$ are marginally trapped. This kind of metrics include the plane waves subset of the “pp-waves” (see e.g. [12, 13, 14, 15] and references therein), although in $D > 4$ they are surprisingly richer as shown in Appendix B of [10]. Still, there is no $\mathcal{H}$ and $S_{t,x}$ are generically non-closed. The case with $G = \text{const.}$ is included here.

3. $G = \sin t \sin x$ with x periodic. These cases allow for topologies S^2 , $\mathbb{R} \times S^1$, etc in the $\{x^a\}$ -part of $(\mathcal{V}, g)$, and are D -generalizations of the Gowdy models [16], including some Robertson-Walker cosmologies. In this case there is a non-trivial $S_{t,x}$ -horizon $\mathcal{H}$, with two connected components, which splits the spacetime into 4 regions, two of them with trapped surfaces $S_{t,x}$ (which may be closed), the other two without them. A similar but open-universe case arises by setting $G = \sinh t \sinh x$.

^Σ Whether or not these surfaces are closed is an open question at this stage, depending on the specific topology of the coordinates $\{x^A\}$.

4. Of course, one can use the general solution $G = f_1(t - x) + f_2(t + x)$, with arbitrary functions f_1, f_2 . This is specially appropriate, with adequate choices of f_1, f_2 , to describe the collision of plane waves (see e.g. [13] and references therein for the GR case; D -generalizations were given in [15, 17]). The standard procedure to build the colliding-wave spacetime is to replace f_1 and f_2 by $f_1(\Theta(t - x)), f_2(\Theta(t + x))$, where Θ is the Heaviside step function. Then, the two regions with $(t - x)(t + x) < 0$ are two plane waves, and the two zones with $(t - x)(t + x) > 0$ correspond to their interaction region and the flat background.

From the above one can derive yet another application. As is known, the previous vacuum solutions can be seen as Kaluza-Klein, or string/M-theory spacetimes which under dimensional reduction become 4-dimensional spacetimes with a number of scalar fields. The scalar fields are given by a subset of the $\{U_A\}$ or appropriate linear combinations of them, see e.g [18, 19]. Actually, there is a (apparently overlooked) one-to-one correspondence between the solutions found in [10] and the solutions generated by the technique explained in section 2 of [19], the $\{U_A\}$ of the former corresponding to the $\{p, \psi_i\}$ of the latter, as can be easily proved. One of the simplest dimensional reductions [18, 19] starts with a line-element of type

$$ds_D^2 = \exp\left(-\sum_{i=4}^{D-1} \psi_i\right) ds_4^2 + \sum_{i=4}^{D-1} e^{2\psi_i} (dx^i)^2$$

where ds_4^2 is a 4-dimensional line-element and $\{x^i\}$ are coordinates on a $(D - 4)$ -torus. As is clear, the physically observable $(D - 2)$ -surfaces are those reduced to 2-surfaces in ds_4^2 . For these, $G^2 = \det g_{AB}$ and $\mathbf{g}_a$ become simply $(A', B', \dots = 2, 3)$

$$G^2 = \exp\left(-2\sum_{i=4}^{D-1} \psi_i\right) (\det g_{A'B'}) \prod_{i=4}^{D-1} e^{2\psi_i} = \det g_{A'B'}$$

$$\mathbf{g}_a = \exp\left(-\sum_{i=4}^{D-1} \psi_i\right) g_{aA'} dx^{A'} \implies g_a^A = \delta_{A'}^A \gamma^{A'B'} g_{aB'}$$

so that the trapping properties of the surfaces S_{X^a} *remain unchanged* whether they are seen as 2-surfaces in $D = 4$ or as $(D - 2)$ -surfaces in full D . Hence, the S_{X^a} -horizon $\mathcal{H}$ is lifted (or reduced!) from 4 to D (D to 4) dimensions. The considered surfaces are closed in D dimensions if and only if they are closed in $D = 4$.

Let us come back to the theoretical approach. The results can be strengthened to the case of *closed* trapped surfaces. Take *any* closed spacelike $(D - 2)$ -surface $\bar{S}$ and adapt the coordinates such that $\bar{S} \subset \Sigma_f \equiv \{f(x^a) = 0\}$ where df is timelike everywhere (apart from this f is arbitrary). This can be done in many different ways. The imbedding (1) for $\bar{S}$ is given by

$$x^a = \Phi^a(\mu^C), \quad x^A = \Phi^A(\mu^C), \quad f(\Phi^a(\mu^C)) = 0, \quad (18)$$

where $\{\mu^C\}$ are intrinsic coordinates on $\bar{S}$. As $\bar{S}$ is compact without boundary, Φ^a must reach their maximum and minimum somewhere on $\bar{S}$. From (18) it follows that $f_{,a} \partial \Phi^a / \partial \mu^C = 0$ so that, at any point $q \in \bar{S}$ where there is an extreme of Φ^1

($\partial\Phi^1/\partial\mu^C|_q = 0$), Φ^0 must have a critical point ($\partial\Phi^0/\partial\mu^C|_q = 0$). This implies, firstly, that the two future-directed null normals of $\bar{S}$ at q are given by $\mathbf{k}^\pm$ of (10), and secondly, that $\det(\partial\Phi^C/\partial\mu^A)|_q \neq 0$ (so that the imbedding (18) has rank $(D-2)$ there.) A way to visualize this is that $\bar{S}$ must be bi-tangent to some S_{X^a} (here the assumption that the surfaces S_{X^a} do not intersect is needed, so that Φ^a are differentiable.) Consequently, we can choose $\{\mu^C\}$ in a neighbourhood $V(q) \subset \bar{S}$ of q such that $\Phi^A(\mu^C) = \mu^A$ at $V(q)$. A straightforward calculation for the traces (6) of $\bar{S}$ leads now to

$$K_{\bar{S}}^\pm|_q = K_{x^a}^\pm|_q - k_a^\pm \gamma_{\bar{S}}^{AB} \frac{\partial^2 \Phi^a}{\partial \mu^A \partial \mu^B} \Big|_q \quad (19)$$

where $K_{x^a}^\pm$ are given in (12) and $\gamma_{\bar{S}}^{AB}$ is the contravariant first fundamental form of $\bar{S}$. The analysis of the second term in the righthand side of (19) can be done as follows. Due to (18)

$$f_{,a} \frac{\partial^2 \Phi^a}{\partial \mu^A \partial \mu^B} \Big|_q = 0 \implies f_{,a} \gamma_{\bar{S}}^{AB} \frac{\partial^2 \Phi^a}{\partial \mu^A \partial \mu^B} \Big|_q = 0,$$

and given that the gradient of f is timelike everywhere, it follows that $\gamma_{\bar{S}}^{AB} \frac{\partial^2 \Phi^a}{\partial \mu^A \partial \mu^B} \Big|_q$ is spacelike with respect to g^{ab} . Hence, as the inverse of g^{ab} is $-2k_{(a}^+ k_{b)}^-$, both $k_a^\pm \gamma_{\bar{S}}^{AB} \frac{\partial^2 \Phi^a}{\partial \mu^A \partial \mu^B} \Big|_q$ must have opposite signs.

Some interesting conclusions can be derived. Assume that H_a is spacelike in a region, then one of the $K_{x^a}^\pm$ is positive and the other is negative for all values of X^a in that region. Letting $K_{x^a}^+ > 0$ (say), then at the maximum or the minimum of Φ^1 one can always show that

$$K_{\bar{S}}^+|_q \geq K_{x^a}^+|_q > 0, \quad K_{\bar{S}}^-|_q \leq K_{x^a}^-|_q < 0$$

and thus the surface $\bar{S}$ is untrapped. In fact, from the above reasoning follows the sufficiency that neither of the $K_{x^a}^\pm$ changes sign. Thus, there are no closed trapped surfaces at the hypersurfaces Σ_f in any region where the S_{X^a} are marginally trapped or absolutely non-trapped.

The last result has many applications too. As a simple but powerful one, consider once again the spherically symmetric line-element (15) and assume that $R_{,\mu}$ is non-timelike everywhere. Then, the $(D-2)$ -spheres are either absolutely non-, or marginally, trapped everywhere. Besides, this implies that *there cannot be any closed trapped surface in $(\mathcal{V}, g)$ at all*. For, due to the structure of the manifold, any closed $\bar{S}$ must be osculating to some $(D-2)$ -sphere somewhere, and at this point their corresponding κ 's of (14) coincide, proving that $\bar{S}$ cannot be trapped there. In particular, *globally* static cases have $R_{,\mu}$ non-timelike everywhere, hence *they cannot contain closed trapped surfaces*. These include flat spacetime, or Einstein's and anti-de Sitter's universes, for example. Of course, this is a known result, but the proof was rather indirect: if there were a closed trapped surface, the spacetime would be geodesically incomplete [1, 2, 3], which is not. It must be noted that any of the mentioned spacetimes, including flat one, certainly contains trapped surfaces (non-compact!, see [3] for examples), so that the previous result is not obvious in principle.

All in all, the significance and applicability of (13-14), which are wholly general, extremely simple, and easily computable, seems worth exploring in detail in the many different gravitational theories. Their potential applications seem to be very many.

Acknowledgement

I thank Roberto Emparan, Alex Feinstein, Marc Mars, and Raúl Vera for their comments and suggestions.

References

- [1] R. Penrose, *Phys. Rev. Lett.* **14**, 57 (1965).
- [2] S. W. Hawking and G. F. R. Ellis, *The large scale structure of space-time*, (Cambridge Univ. Press, Cambridge, 1973).
- [3] J.M.M. Senovilla, *Gen. Rel. Grav.* **30**, 701 (1998).
- [4] M. Kriele, *Spacetime*, (Springer, Berlin, 1999).
- [5] F.R. Tangherlini, *Nuovo Cimento* **27**, 636 (1963).
- [6] S.A. Hayward, *Phys. Rev. D* **53**, 1938 (1996).
- [7] J.F. Villas da Rocha and A. Wang, *Class. Quantum Grav.* **17**, 2589 (2000).
- [8] R. Emparan and H.S. Reall, *Phys. Rev. Lett.* **88**, 101101 (2002).
- [9] R.C. Myers and M. J. Perry, *Ann. Phys. (N.Y.)* **172**, 304 (1986).
- [10] R. Emparan and H.S. Reall, *Phys. Rev. D* **65**, 084025 (2002).
- [11] V. D. Ivashchuk and V. N. Melnikov, *Grav.Cosmol.* **1**, 204 (1995).
- [12] D. Kramer *et al.*, *Exact Solutions of Einstein's Field Equations*, (Cambridge Univ. Press, Cambridge, 1980).
- [13] J. B. Griffiths, *Colliding plane waves in General Relativity*, (Oxford University Press, Oxford, 1991).
- [14] C. M. Hull, *Phys. Lett. B* **139**, 39 (1984); J.M. Figueroa-O'Farrill, *Class. Quantum Grav.* **17**, 2925 (2000); R. Gueven, *Phys. Lett. B* **482**, 255 (2000).
- [15] P.M. Schwarz, *Nuc. Phys. B* **373**, 529 (1992).
- [16] R.H. Gowdy, *Phys. Rev. Lett.* **27**, 286 (1971).
- [17] V. Bozza and G. Veneziano, *JHEP* **0010**, 035 (2000); A. Feinstein, K. Kunze and M.A.Vázquez-Mozo, *Class. Quantum Grav.* **17**, 3599 (2000).
- [18] H.Lü and C.N. Pope, *Nuc. Phys. B* **465**, 127 (1996); N.A. Obers and B. Pioline, *Phys. Rep.* **318**, 113 (1999).
- [19] A. Feinstein and M.A.Vázquez-Mozo, *Nuc. Phys. B* **568**, 405 (2000).